

REPLY TO GLOBAL HIGH-ALTITUDE LIMITS FOR AMPHIBIANS BY TRACIE A. SEIMON AND ANTON SEIMON (2015)

Barkha Subba¹, G. Ravikanth² & N.A. Aravind³

^{1,2,3} Ashoka Trust for Research in Ecology and the Environment (ATREE), Suri Sehgal Centre for Biodiversity and Conservation, Royal Enclave, Srirampura, Jakkur PO, Bengaluru, Karnataka 560064, India

¹ Manipal University, Manipal, Karnataka 576104, India

¹ barkhasubba@gmail.com (corresponding author),

² gravikanth@atree.org, ³ aravind@atree.org

High altitude records of amphibians in the age of Climate Change

With the problem of global warming looming large on our planet the existence of many species are under threat. Researchers are racing to bring in new information on the impact of climate change on global biodiversity (Walther et al. 2002; Parmesan & Yohe 2003; Campbell et al. 2009; Xu et al. 2009; Bellard et al. 2012; Gottfried et al. 2012; Shrestha et al. 2012). There have been a large number of studies that have documented range shifts of species due to climate change (Walther et al. 2005; Araujo & Rahbek 2006; Hickling et al. 2006; Chen et al. 2011). The new records on altitude of both flora and fauna as a result of range extension due to climate change have become quite frequent. In this context, we agree with Seimon et al. (2015) who have neatly summarized the three problems on new findings about altitude records in their Response to our paper (Subba et al. 2015). Furthermore, we agree with authors that often vital data may be hidden within other larger issues; as a result missing out on relevant data seems to be a reoccurring problem (Seimon et al. 2007a). In this age of easy access to information and an era where scientists and citizens are coming together to bridge the gap in information, there is a need to make the data easily available and visible. New records

should also be followed by the exact location (latitude - longitude) where the species was observed or collected along with the stage of development.

In their comment Seimon et al. mention that we did not take into account the record of 5200m for *Scutigera alticola (boulengeri)* in Tibet recorded by Swan (1990). This was an intentional omission, the reason being the following—

Swan (1990) writes “*The Himalayan and Tibetan frogs of the Family Pelobatidae include Scutigera sikkimensis that is found in scattered localities along the Himalaya between 2800 and 4000 m and Scutigera alticola which is primarily Tibetan and has been collected up to 5200m.*”

Here he fails to mention who collected it, he may have been referring to Hocks record back in 1964, and we surely know that Swan did not make the collection from the following lines, where he writes, “*In 1980 I was privileged to accompany the first international tour across Tibet as a guest of the Chinese Academy of Sciences where I was able to see and collect in places where the fabled explorers of the last century made their remarkable observations. On a pass (Gyaco la, northeast of Tingri) I noticed some snow fed ponds and was drawn to them hoping for specimens of Scutigera alticola but the black mud I scooped up finally revealed two small scaleless loaches (Nemachilinae). This was at 5200 m and somewhat higher than similar fish obtained by G.E. Hutchinson in Ladakh back in 1932 and so they represented the highest fish in the world.*”

We opine that not only new altitudinal records but old records as well require screening to some extent.

In our paper we did not write about other high altitude Himalayan frogs, as it was not within the scope of this short communication paper as the article is not exclusively about altitudinal distribution of amphibians.



ISSN 0974-7907 (Online)
ISSN 0974-7893 (Print)

OPEN ACCESS



DOI: <http://dx.doi.org/10.11609/JoTT.o4405.7851-2>

Date of publication: 26 September 2015 (online & print)

Manuscript details: Ms # o4405 | Received 10 September 2015

Citation: Subba, B., G. Ravikanth & N.A. Aravind (2015). Reply to global high-altitude limits for amphibians by Tracie A. Seimon and Anton Seimon (2015). *Journal of Threatened Taxa* 7(11): 7851–7852; <http://dx.doi.org/10.11609/JoTT.o4405.7851-2>

Copyright: © Subba et al. 2015. Creative Commons Attribution 4.0 International License. JoTT allows unrestricted use of this article in any medium, reproduction and distribution by providing adequate credit to the authors and the source of publication.

We have mainly focused on *S. boulengeri* and *S. sikimensis*, the reason being, *S. boulengeri* is often misidentified as *S. sikimensis*, and this information would be very useful for future researchers in this area.

We acknowledge that we missed the paper by Seimon et al. (2007b) on range extension of Andean anurans. The authors have recorded *Pleuroderma* sp. at a deglaciated seasonal pool at an altitude of 5400m. This is a very important and interesting paper on range extension by relatively lower altitude species to higher altitudes due to deglaciation. At the same time, we would like to point out how the well known and widely read book *Ecology and Behaviour of Amphibians* by Kentwood (Wells 2007) and reputable websites like AmphibiaWeb (<http://amphibiaweb.org/>. Accessed: 1.9.2015) also unfortunately fails to mention Seimon et al.'s altitude record of *Pleuroderma* sp. on their section on altitudinal distribution of amphibians in the world. They have mentioned *S. boulengeri* as probably the highest altitude amphibian in the world. We are writing this in hope that publication on altitude record gets more visibility as it has become a vital element in studying species response to the rapidly changing climate. Regarding the ephemeral pool where the authors photographed the tadpoles of *Pleuroderma* sp., the location would have been much more clear if the authors had given exact point location (latitude longitude) along with altitude (5400m), for the sake of "consistency and precision" of altitudinal records.

Lake Gurudongmar Complex 2, one of the lakes where we found *S. boulengeri* has increased in size from 0.249km² in 1965 to 1.076km² in 2010 (Kumar & Prabhu 2012) as a result of active deglaciation. In its home range in the eastern Himalaya, the highest authentic record of *Scutigera boulengeri*, with point location of the collection made, is at an altitude of 4324m (Chen et al. 2009; Li et al. 2009) and that of an anecdotal and unsubstantiated record is 5200m (Swan 1990). We collected an adult specimen from among a healthy population of *S. boulengeri* at an altitude of 5270m at Lake Gurudongmar Complex 2. This Lake Complex, including the seasonal pools in its vicinity has an altitudinal range of 5250–5490m. We reaffirm that *S. boulengeri* is undergoing upward range expansion and indeed "scaling new heights."

References

- Araújo, M.B. & C. Rahbek (2006). How does climate change affect biodiversity?. *Science - New York Then Washington* 313(5792): 1396.
- Bellard, C., C. Bertelsmeier, P. Leadley, W. Thuiller & F. Courchamp (2012). Impacts of climate change on the future of biodiversity. *Ecology letters* 15(4): 365–377; <http://dx.doi.org/10.1111/j.1461-0248.2011.01736.x>
- Campbell, A., V. Kapos, J.P. W. Scharlemann, P. Bubb, A. Chenery, L. Coad, B. Dickson, N. Doswald, M.S.I. Khan, F. Kershaw & M. Rashid (2009). Review of the Literature on the Links between Biodiversity and Climate Change: Impacts, Adaptation and Mitigation. Secretariat of the Convention on Biological Diversity, Montreal. Technical Series No. 42, 124pp.
- Chen, W., K. Bi & J. Fu (2009). Frequent mitochondrial gene introgression among high elevation Tibetan megophryd frogs revealed by conflicting gene genealogies. *Molecular Ecology* 18(13): 2856–2876; <http://dx.doi.org/10.1111/j.1365-294X.2009.04258.x>
- Chen, I.C., J.K. Hill, R. Ohlemüller, D.B. Roy & C.D. Thomas (2011). Rapid range shifts of species associated with high levels of climate warming. *Science* 333(6045): 1024–1026.
- Gottfried, M., H. Pauli, A. Futschik, M. Akhalkatsi, P. Barančok, J.L.B. Alonso & G. Grabherr (2012). Continent-wide response of mountain vegetation to climate change. *Nature Climate Change* 2(2): 111–115.
- Hickling, R., D.B. Roy, J.K. Hill, R. Fox & C.D. Thomas (2006). The distributions of a wide range of taxonomic groups are expanding polewards. *Global Change Biology* 12(3): 450–455.
- Hock, R.J. (1964). Animals in high altitudes: reptiles and amphibians, pp. 841–842. In: Dill, D.B., E.F. Adolph & C.G. Wilber (eds.). *Handbook of Physiology, Sect 4, Adaptation to Environment*. American Physiological Society, Washington DC, 1056pp.
- Kumar, B. & T.S. Prabhu (2012). Impact of Climate Change: Glacier Lake Outburst Floods (GLOFs), pp. 81–102. In: Arrawatia, M.L. & S. Tambe (eds.). *Climate Change in Sikkim - Patterns, Impacts and Initiatives*. Information and Public Relations Department, Government of Sikkim, xxiii +424pp.
- Li, R., W. Chen, L. Tu & J. Fu (2009). Rivers as barriers for high elevation amphibians: a phylogeographic analysis of the alpine stream frog of the Hengduan Mountains. *Journal of Zoology* 277: 309–316; <http://dx.doi.org/10.1111/j.1469-7998.2008.00543.x>
- Parmesan, C. & G. Yohe (2003). A globally coherent fingerprint of climate change impacts across natural systems. *Nature* 421: 37–42.
- Seimon, A., S.R.P. Halloy & T.A. Seimon (2007a). Recent observation of a proliferation of *Ranunculus trichophyllus* Chaix. in high-altitude lakes of the Mount Everest region: Comment. *Arctic, Antarctic and Alpine Research* 39: 340–341.
- Seimon, T., A. Seimon, P. Daszak, S.R.P. Halloy, L.M. Schloegel, C.S. Aguilar, P. Sowell, A.D. Hyatt, B. Konecky & J.E. Simmons (2007b). Upward range extension of Andean anurans and chytridiomycosis to extreme elevations in response to tropical deglaciation. *Global Change Biology* 13: 288–299.
- Shrestha, U.B., S. Gautam & K.S. Bawa (2012). Widespread climate change in the Himalayas and associated changes in local ecosystems. *PLoS ONE* 7(5): e36741; <http://dx.doi.org/10.1371/journal.pone.0036741>
- Subba, B., G. Ravikanth & N.A. Aravind (2015). Scaling new heights: first record of Boulenger's Lazy Toad *Scutigera boulengeri* (Amphibia: Anura: Megophryidae) from high altitude lake in Sikkim Himalaya, India. *Journal of Threatened Taxa* 7(10): 7655–7663; <http://dx.doi.org/10.11609/JoTT.o4325.7655-63>
- Swan, L. (1990). The highest life. *Himalayan Journal* 46: 125–133.
- Walther, G.R., E. Post, P. Convey, A. Menzel, C. Parmesan, J.C.B. Tervey, J.M. Formentin, O.H. Guldberg & F. Bairlein (2002). Ecological responses to recent climate change. *Nature* 416: 389–395.
- Walther, G.R., S. Beißner, & C.A. Burga (2005). Trends in the upward shift of alpine plants. *Journal of Vegetation Science* 16(5): 541–548.
- Wells, K.D. (2007). *Ecology and Behaviour of Amphibians*. University of Chicago Press, xvi+1148pp.
- Xu, J., R.E. Grumbine, A. Shrestha, M. Eriksson, X. Yang, Y. Wang, & A. Wilkes (2009). The Melting Himalayas: Cascading Effects of Climate Change on Water, Biodiversity, and Livelihoods. *Conservation Biology* 23: 520–530; <http://dx.doi.org/10.1111/j.1523-1739.2009.01237.x>
- AmphibiaWeb Information on amphibian biology and conservation. [web application] 2015. Berkeley, California: AmphibiaWeb. Available:<http://amphibiaweb.org/>. (Accessed: 01 September 2015)

